

A NOTE ON THE TRIVIALITY OF κ -DEFORMATIONS OF GRAVITY

D. GRUMILLER, W. KUMMER, D.V. VASSILEVICH¹

УДК 11.10.Lm; 11.80.Et
© 2001 p.

Theoretische Physik, Technische Universität Wien
(Wiedner Hauptstr. 8-10, A-1040 Wien, Austria)

¹ Institut für Theoretische Physik, Universität Leipzig
(Augustusplatz 10, D-04109 Leipzig, Germany)

Some simple observations concerning certain κ -deformations in the framework of 2D dilaton gravity formulated as Poisson-sigma models are discussed, and the question of (non-)triviality is addressed.

1. Introduction

Certain astrophysical observations have motivated the study of κ -deformed Poincaré-algebras, e.g. in the context of “Doubly Special Relativity” (henceforth DSR; for a recent review and more references cf. e.g. [1, 2, 3]). It might be helpful to consider the simpler case of $D = 2$ first to settle open conceptual questions (especially concerning gravity). The first work in this direction appeared only recently [4]. In the present note when talking about κ -deformations¹ we will exclusively refer to the deformation introduced by Magueijo and Smolin [3], which differs from the original proposal [5].²

Moreover, it seems to be an interesting task by itself for purely mathematical reasons. Indeed, Izawa has shown a few years ago, that the most general consistent deformation (in the sense of Barnich and Henneaux [7]) of a Poisson- σ model (PSM) is again a PSM with the same number of target space coordinates [8]. Since dilaton gravity without matter in 2D is merely a very special PSM [9] this result applies to it as well. Thus, generic consistent deformations of dilaton gravity in 2D are mathematically feasible.

The description of dilaton gravity in terms of a PSM has turned out to be very fruitful – e.g. all classical solutions have been obtained locally and globally within this approach [10] and even in the presence of matter an exact quantization of geometry has been achieved

[11], with interesting phenomenological applications for scattering processes [12]. For a recent review on dilaton gravity in 2D ref. [13] can be consulted.

In the present note we intend to discuss a very simple question in the framework of twodimensional dilaton gravity: are κ -deformations trivial or not (and in what sense are they (non)trivial)?

2. Triviality of deformations

Let the momenta p_a and the generator of boosts J satisfy the undeformed Poincaré algebra in 2D:

$$[p_0, p_1] = 0, \quad [J, p_0] = p_1, \quad [J, p_1] = p_0. \quad (1)$$

An elementary calculation shows that the generators

$$P_a = \frac{p_a}{1 + p_0/\kappa}, \quad J, \quad (2)$$

with $a = 0, 1$ satisfy the κ -deformed relations:

$$\begin{aligned} [P_a, P_b] &= 0, & [J, P_0] &= P_1 - \kappa^{-1} P_1 P_0, \\ [J, J] &= 0, & [J, P_1] &= P_0 - \kappa^{-1} P_1^2. \end{aligned} \quad (3)$$

Obviously for $\kappa \rightarrow \infty$ the undeformed algebra is recovered. It is also instructive to invert (2):

$$p_a = \frac{P_a}{1 - P_0/\kappa}. \quad (4)$$

Clearly the deformation is algebraically trivial.³ The whole effect is a change of variables (2). Therefore, it seems natural to do the *same* change of variables in the PSM, the action of which reads [9]

$$L_{\text{PSM}} = \int_{M_2} \left[dX^I \wedge A_I + \frac{1}{2} \mathcal{P}^{IJ} A_J \wedge A_I \right]. \quad (5)$$

¹Of course, κ -deformations do not automatically imply DSR, nor vice versa.

²By κ -deformation of Poincaré algebra only the deformation of the algebraic sector is considered in the present work, characterized by some fundamental mass parameter κ . In [5] there was proposed the framework with quantum κ deformation, introducing as well the deformed nonsymmetric coproducts in the coalgebra sector, which leads to noncommutative space-time [6].

³This has been pointed out in ref. [2]. This statement also reminds one of the rigidity results on deformations of Lie algebras [14].

For dilaton gravity (assuming Lorentzian signature for definiteness) the three gauge field 1-forms are $A_I = (\omega, e^-, e^+)$, where $e^\pm$ is the dual basis of 1-forms in light-cone gauge for the anholonomic frame and ω is related to the spin-connection 1-form via $\omega^a_b = \varepsilon^a_b \omega$. The three target space coordinates $X^I = (X, X^+, X^-)$ contain the so-called “dilaton” X and two auxiliary fields $X^\pm = (X^0 \pm X^1)$ (again in light-cone representation). The Poisson-tensor $\mathcal{P}^{IJ}$ depends solely on the target space coordinates, it is antisymmetric and fulfills the Jacobi-identity $\mathcal{P}^{IN}\partial_N\mathcal{P}^{JK} + \text{cycl.}(IJK) = 0$. Due to the odd number of target space coordinates this tensor cannot have full rank and hence for non-trivial models exactly one Casimir function exists, related to the “energy” of the spacetime [15].

The simplest dilaton gravity model is obtained with the “free” Poisson tensor

$$\mathcal{P} = \begin{pmatrix} 0 & 0 & -X^1 \\ 0 & 0 & -X^0 \\ X^1 & X^0 & 0 \end{pmatrix}, \quad (6)$$

yielding vanishing curvature and torsion. In primed coordinates

$$X^{a'} = \frac{X^a}{1 + X^0/\kappa}, \quad X' = X \quad (7)$$

the Poisson tensor transforms as

$$\mathcal{P}^{I'J'} = \frac{\partial X^{I'}}{\partial X^I} \frac{\partial X^{J'}}{\partial X^J} \mathcal{P}^{IJ}. \quad (8)$$

Consequently, in the primed coordinates we obtain again the free PSM with an extra term,

$$\kappa^{-1} X^{1'} X^{a'} \omega \wedge e^{a'}, \quad (9)$$

which is the result of Mignemi [4].

It seems natural to deform other dilaton models (spherically reduced gravity, the CGHS model [16], the JT model [17], etc.) in the same way [18] and to study the corresponding global structure. Thus, the starting point is the (undeformed) dilaton gravity action

$$L = \int_{M_2} [X_a D e^a + X d\omega + \epsilon (X^+ X^- U(X) + V(X))] , \quad (10)$$

with $\epsilon = e^+ \wedge e^-$ being the volume 2-form and $X_a D e^a = X^+(d - \omega) \wedge e^- + X^-(d + \omega) \wedge e^+$ contains the torsion 2-form in light-cone representation. It is actually not necessary to restrict oneself to potentials of the type $X^+ X^- U(X) + V(X)$ – a particular class of relevant counter examples can be found in ref. [19] – but

for sake of simplicity this special form will be assumed in the present work because it covers all special models referred to above. The following equations will be useful

$$\begin{aligned} \frac{\partial X^{0'}}{\partial X^0} &= \left(1 - \frac{X^{0'}}{\kappa}\right)^2, & \frac{\partial X^{1'}}{\partial X^1} &= 1 - \frac{X^{0'}}{\kappa}, \\ \frac{\partial X^{1'}}{\partial X^0} &= -\frac{X^{1'}}{\kappa} \left(1 - \frac{X^{0'}}{\kappa}\right). \end{aligned} \quad (11)$$

By employing (8), (11) together with the relation

$$X^+ X^- = \frac{X^{+'} X^{-'}}{\left(1 - \frac{X^{0'}}{\kappa}\right)^2}, \quad (12)$$

the transformation law for the potentials is established:

$$U \rightarrow U' = U \left(1 - \frac{X^{0'}}{\kappa}\right), \quad V \rightarrow V' = V \left(1 - \frac{X^{0'}}{\kappa}\right)^3 \quad (13)$$

Of course, the change of variables considered above is not an equivalence transformation: (i) it is singular; (ii) it changes the definition of the metric.

Invariants of the deformed model can be traced back easily to the undeformed case. For example, the “line element”

$$\begin{aligned} (ds)^2 &= e^{0'} \otimes e^{0'} \left(1 - \frac{\eta_0}{\kappa}\right)^4 - (e^{1'} \otimes e^{0'} + e^{0'} \otimes e^{1'}) \frac{\eta_1}{\kappa} \times \\ &\times \left(1 - \frac{\eta_0}{\kappa}\right)^3 - e^{1'} \otimes e^{1'} \left(1 - \frac{\eta_0}{\kappa}\right)^2 \left(1 - \frac{\eta_1^2}{\kappa^2}\right) \end{aligned} \quad (14)$$

with $\eta_a := X^{a'}$ results from a target space diffeomorphism (8) of the PSM, applied to the standard $(ds)^2$ and therefore must be invariant under the deformed transformations. Clearly, a description of the deformed geometry by means of (14) does not make much sense because the deformation would be without effect on classical solutions since all singularities in $e^{0'}, e^{1'}$ are compensated by corresponding zeros. This follows already from the previous general considerations, but it can be checked explicitly by plugging the explicit solutions (15) below into (14). The result is the original line element $(ds)^2 = e^0 \otimes e^0 - e^1 \otimes e^1$.

On the other hand, taking a non-invariant object as “metric” seems to be questionable from a physical point of view. For instance, the curvature scalar related to the metric presented in [4, 18] depends on the choice of gauge, which from a relativistic point of view is an inconvenient feature.

Thus there seems to be no way to evade both, the Scylla of triviality and the Charybdis of non-invariance.

3. Deformed solutions

For a more explicit analysis we have to express the deformed “zweibein” in terms of the undeformed variables,

$$\begin{aligned} e^{0'} &= \left(1 + \frac{X^0}{\kappa}\right)^2 e^0 + \frac{X^1}{\kappa} \left(1 + \frac{X^0}{\kappa}\right) e^1, \\ e^{1'} &= \left(1 + \frac{X^0}{\kappa}\right) e^1. \end{aligned} \quad (15)$$

If we require non-triviality of the deformation a non-invariant “metric” has to be employed, as discussed in the previous section. Since there does not seem to exist a better alternative (there is no canonical choice), let us assume that the “physical metric” is constructed in the usual way from the deformed “zweibein” (15), namely $g'_{\mu\nu} = e_{\mu}^{a'} \otimes e_{\nu}^{b'} \eta_{ab}$, where η_{ab} is the flat metric.⁴

As a demonstration one can choose spherically reduced gravity (from arbitrary dimension D) as undeformed starting point. Equations of motion imply

$$X^a = \epsilon^{\mu\nu} e_{\mu}^a \partial_{\nu} X. \quad (16)$$

The discussion will be restricted to the behavior of the deformed solution in the asymptotic region of the undeformed one. This is the simplest test for the validity of the deformed model. In this region we can apply Schwarzschild gauge $(ds)^2 = \xi(dt)^2 - \xi^{-1}(dr)^2$ with $\xi(r)$ and $X(r)$ defined by [20]

$$\xi(r) = 1 + 2\mathcal{C}_0 |1 - a|^{\frac{a}{a-1}} r^{\frac{a}{a-1}} \left(\frac{B}{a}\right)^{\frac{2-a}{2(a-1)}}, \quad (17)$$

and

$$r = \sqrt{\frac{a}{B}} \frac{1}{|1 - a|} X^{1-a}, \quad (18)$$

respectively. The constants in these equations have the following meaning: $a = (D - 3)/(D - 2)$, B is a normalization constant (which can be set to 1) and $\mathcal{C}_0$ is the value of the Casimir function (for positive/negative values a naked singularity/black hole is described, respectively; if it vanishes the Minkowskian ground state is reached). The (singular) limit $D \rightarrow \infty$, if treated with care, yields the CGHS model.

Asymptotically ($r \rightarrow \infty$) one readily obtains

$$X \sim r^{1/(1-a)}, \quad X^1 = 0, \quad X^0 \sim r^{a/(1-a)}. \quad (19)$$

⁴The words “metric” and “zweibein” have been put under quotation marks to indicate that this nomenclature is too suggestive. Neither $e^{a'}$ nor $g'_{\mu\nu}$ transform as the notation seems to promise (unless κ is taken to ∞).

⁵It is somewhat amusing that $D = 4$ is the limiting case.

One observes that for $\frac{1}{2} \leq a < 1$ (i.e. for spherical reduction with $4 \leq D < \infty$) the function

$$\chi(r) = 1 + \frac{X^0}{\kappa} \quad (20)$$

which characterizes the “strength” of the deformation is large, implying that asymptotically the model is being deformed noticeably.⁵ This indicates that we are dealing in fact with the old Fock version [21] of the deformations which modifies physics at large distances. Indeed, the way in which the generators X^I have been interpreted corresponds to a deformation in coordinate space rather than in momentum space: for instance, the dilaton X has been regarded as some (power of) a “radius” and thus as a coordinate space entity. However, these interpretational issues are not pivotal for the present discussion.

Despite of this sizable asymptotic deformation the scalar curvature for the primed zweibein reads

$$R(r) \sim r^{-2/(1-a)}. \quad (21)$$

Thus, the asymptotically flat region remains asymptotically flat after the deformation, which is an attractive feature and indicates that it may be sensible to talk of an “asymptotic region” even in the deformed case.

It should be emphasized strongly, though, that one should not over-interpret results extracted from a “metric” which does not possess the (deformed) Lorentz invariance.

4. Conclusion

We have shown that κ -deformed dilaton gravity in $2D$ is either classically equivalent to a corresponding undeformed model (and thus the deformation would be trivial) or one has to deal with a “non-invariant metric”.

However, this does not imply that all deformations are without effect. In particular, the question of the “correct” metric turns out to be a non-trivial one, even after imposing the invariance condition. This will be the subject of work in progress dealing with these issues in a much more comprehensive manner [22, 23].

We stress that only the classical part of the full κ -deformed Poincaré bialgebra is being used in our construction. Our procedure may be as well called “deformed gravity with an invariant energy scale”. An interesting development may consist in a combination of the

Poisson- σ models with the quantum algebra approach to DSR described in the recent papers [24].

It also would be worthwhile to investigate to which extent our conclusions depend on a particular choice of the basis in the κ -deformed Poincaré algebra.

During the final preparations of this proceedings contribution an e-print appeared which has partial overlap with our discussion [25]. The authors of that paper conclude that DSR is operationally indistinguishable from special relativity.

Acknowledgement

We would like to thank our collaborators on deformations of dilaton gravity in 2D, S. Mignemi and T. Strobl, for enlightening discussions and helpful correspondence. Moreover, we are indebted to V. Lyakhovsky for his comments on deformations of algebras. We thank G. Amelino-Camelia for his interest and correspondence. Finally, we are grateful to J. Lukierski for relevant critical remarks.

One of the authors (D.G.), who has presented the results of this note at the XIVth International Hutsulian Workshop on “Mathematical Theories and their Physical & Technical Applications” in Cernivtsi (Ukraine), is grateful to the organizers and participants of this meeting for their hospitality and numerous interesting conversations, in particular L. Adamska, S. Moskaliuk, S. Rumyantsev, A. Vlassov and M. Wohlgenannt.

This work has been supported by project P-14650-TPH of the Austrian Science Foundation (FWF), by DFG project BO 1112/12-1, by the MPI MIS (Leipzig) and by the Austro-Ukrainian Institute for Science and Technology.

1. *Amelino-Camelia G.* Doubly-special relativity: First results and key open problems // arXiv.org/abs/gr-qc/0210063.
2. *Maguiejo J. and Smolin L.* Generalized Lorentz invariance with an invariant energy scale // http: // arXiv.org/abs/gr-qc/0207085.
3. *Maguiejo J. and Smolin L.* Lorentz invariance with an invariant energy scale // Phys. Rev. Lett. – 2002. – **88**. – 190403 // arXiv.org/abs/hep-th/0112090.
4. *Mignemi S.* Two-dimensional gravity with an invariant energy scale // arXiv.org/abs/hep-th/0208062.
5. *Lukierski J., Ruegg H., Nowicki A. and Tolstoi V. N.* Q -deformation of Poincaré algebra // Phys. Lett. – 1991. – **B264**. – 331.
6. *S. Majid and H. Ruegg* Bicrossproduct structure of kappa Poincaré group and noncommutative geometry // Phys. Lett. – 1994. – **B334**. – 348 // arXiv.org/abs/hep-th/9405107.
J. Lukierski, H. Ruegg and W. J. Zakrzewski Classical quantum mechanics of free kappa relativistic systems // Annals Phys. – 1995. – **243**. – 90 // arXiv.org/abs/hep-th/9312153.
7. *Barnich G. and Henneaux M.* Consistent couplings between fields with a gauge freedom and deformations of the master equation // Phys. Lett. – 1993. – **B311**. – 123–129 // arXiv.org/abs/hep-th/9304057;
Barnich G., Brandt F., and Henneaux M. Local BRST cohomology in gauge theories // Phys. Rept. – 2000. – **338**. – 439–569 // arXiv.org/abs/hep-th/0002245.
8. *Izawa K. I.* On nonlinear gauge theory from a deformation theory perspective // Prog. Theor. Phys. – 2000. – **103**. – 225–228 // arXiv.org/abs/hep-th/9910133.
9. *Schaller P. and Strobl T.* Poisson structure induced (topological) field theories // Mod. Phys. Lett. – 1994. – **A9**. – 3129–3136 // arXiv.org/abs/hep-th/9405110.
10. *Klösch T. and Strobl T.* Classical and quantum gravity in (1+1)-dimensions. Part 1: A unifying approach // Class. Quant. Grav. – 1996. – **13**. – 965–984 // arXiv.org/abs/gr-qc/9508020;
Klösch T. and Strobl T. Classical and quantum gravity in 1+1 dimensions. Part II: The universal coverings // Class. Quant. Grav. – 1996. – **13**. – 2395–2422 // arXiv.org/abs/gr-qc/9511081;
Klösch T. and Strobl T. Classical and quantum gravity in 1+1 dimensions. Part III: Solutions of arbitrary topology and kinks in 1+1 gravity // Class. Quant. Grav. – 1997. – **14**. – 1689–1723 // arXiv.org/abs/hep-th/9607226;
Klösch T. and Strobl T. A global view of kinks in 1+1 gravity // Phys. Rev. – 1998. – **D57**. – 1034 // arXiv.org/abs/gr-qc/9707053;
Strobl T. Gravity in two spacetime dimensions // arXiv.org/abs/hep-th/0011240.
11. *Kummer W., Liebl H., and Vassilevich D. V.* Exact path integral quantization of generic 2+ d dilaton gravity // Nucl. Phys. – 1997. – **B493**. – 491–502 // arXiv.org/abs/gr-qc/9612012;
Integrating geometry in general 2d dilaton gravity with matter // Nucl. Phys. 1999. – **B544**. – 403–431 // arXiv.org/abs/hep-th/9809168; *Grumiller D.* Quantum dilaton gravity in two dimensions with matter. // PhD thesis, Technische Universität Wien, 2001 // arXiv.org/abs/gr-qc/0105078.
12. *Grumiller D., Kummer W., and Vassilevich D. V.* The virtual black hole in 2d quantum gravity // Nucl. Phys. – 2002. – **B580**. – 438–456 // arXiv.org/abs/gr-qc/0001038;
Fischer P., Grumiller D., Kummer W., and Vassilevich D. V. S -matrix for s -wave gravitational scattering // Phys. Lett. – 2001. – **B521**. – 357–363 // arXiv.org/abs/gr-qc/0105034.
Erratum *ibid.* – 2002. – **B532**. – 373;
Grumiller D. The virtual black hole in 2D quantum gravity and its relevance for the S -matrix // Int. J. Mod. Phys. – 2001. – **A 17**. – 989 // arXiv.org/abs/hep-th/0111138;
Virtual black hole phenomenology from 2d dilaton theories // Class. Quant. Grav. – 2002. – **19**. – 997–1009 // arXiv.org/abs/gr-qc/0111097;
Grumiller D., Kummer W., and Vassilevich D. V. Virtual black holes in generalized dilaton theories and their special role in string gravity // arXiv.org/abs/hep-th/0208052.
13. *Grumiller D., Kummer W., and Vassilevich D. V.* Dilaton gravity in two dimensions // Phys. Rept. – 2002. – **369**. – 327–429 // arXiv.org/abs/hep-th/0204253.
14. *Gerstenhaber M.* On the deformation of rings and algebras // Ann. Math. – 1964. – **79**. – 59;
Levy-Nahas M., Deformation and contraction of Lie algebras

- // J. Math. Phys. – 1967. – **8**. – 1211;
Lyakhovsky V. D. Generalized symmetries and deformations of the direct sums of Lie algebras // Commun. Math. Phys. – 1968. – **11**. – 131.
15. *Grosse H., Kummer W., Presnajder P., and Schwarz D. J.* Novel symmetry of nonEinsteinian gravity in two dimensions // J. Math. Phys. – 1992. – **33**. – 3892–3900 // arXiv.org/abs/hep-th/9205071;
Mann R. B., Conservation laws and $2-d$ black holes in dilaton gravity // Phys. Rev. – 1993. – **D47**. – 4438–4442 // arXiv.org/abs/hep-th/9206044;
Kummer W., and Widerin P. Non-Einsteinian gravity in $d = 2$: Symmetry and current algebra // Mod. Phys. Lett. – 1994. – **A9**. – 1407–1414;
 Conserved quasilocal quantities and general covariant theories in two-dimensions // Phys. Rev. – 1995. – **D52**. – 6965–6975 // arXiv.org/abs/gr-qc/9502031;
Kummer W., and Lau S. R. Boundary conditions and quasilocal energy in the canonical formulation of all $1 + 1$ models of gravity // Annals Phys. – 1997. – **258**. – 37–80 // arXiv.org/abs/gr-qc/9612021;
Kummer W. and Tieber G. Universal conservation law and modified Nöther symmetry in $2d$ models of gravity with matter // Phys. Rev. – 1999. – **D59**. – 044001 // arXiv.org/abs/hep-th/9807122;
Grumiller D. and Kummer W. Absolute conservation law for black holes // Phys. Rev. – 2000. – **D61**. – 064006 // arXiv.org/abs/gr-qc/9902074.
 16. *Callan C. G. (Jr.), Giddings S. B., Harvey J. A., and Strominger A.* Evanescent black holes // Phys. Rev. – 1992. – **D45**. – 1005–1009 // arXiv.org/abs/hep-th/9111056.
 17. *Barbashov B. M., Nesterenko V. V., and Chervyakov A. M.* The solitons in some geometrical field theories // Theor. Math. Phys. – 1979. – **40**. – 572–581 (Teor. Mat. Fiz. – 1979. – **40**. – 15–27), J. Phys. – 1980. – **A13**. – 301–312;
D'Hoker E., and Jackiw R. Liouville field theory // Phys. Rev. – 1982. – **D26**. – 3517;
Teitelboim C. Gravitation and Hamiltonian structure in two space-time dimensions // Phys. Lett. – 1983. – **B126**. – 41;
D'Hoker E., Freedman D., and Jackiw R. $SO(2,1)$ invariant quantization of the Liouville theory // Phys. Rev. – 1983. – **D28**. – 2583;
D'Hoker E., and Jackiw R. Space translation breaking and compactification in the Liouville theory // Phys. Rev. Lett. – 1983. – **50**. – 1719–1722;
Jackiw R. Another view on massless matter-gravity fields in two-dimensions // arXiv.org/abs/hep-th/9501016.
 18. *Mignemi S.* Two-dimensional gravity with an invariant energy scale and arbitrary dilaton potential // arXiv.org/abs/hep-th/0210213.
 19. *Grumiller D. and Vassilevich D. V.* Non-existence of a dilaton gravity action for the exact string black hole // JHEP. – 2002. – **11**. – 018 // arXiv.org/abs/hep-th/0210060.
 20. *Katanaev M. O., Kummer W., and Liebl H.* On the completeness of the black hole singularity in $2d$ dilaton theories // Nucl. Phys. – 1997. – **B486**. – 353–370 // arXiv.org/abs/gr-qc/9602040.
 21. *Fock V. A.* The theory of space, time and gravitation. – Oxford: Pergamon Press, 1964;
Manida S. N. Fock-Lorentz transformations and time-varying speed of light // arXiv.org/abs/gr-qc/9905046.
 22. *Strobl T.* (in preparation).
 23. *Grumiller D., Kummer W., Mignemi S., Strobl T., and Vassilevich D.* Deformations of dilaton gravity in $2d$ (in preparation).
 24. *Kowalski-Glikman J., and Nowak S.* Non-commutative space-time of doubly special relativity theories // arXiv.org/abs/hep-th/0204245;
Lukierski J., and Nowicki A. Nonlinear and quantum origin of doubly infinite family of modified addition laws for four-momenta // arXiv.org/abs/hep-th/0209017;
Kowalski-Glikman J. Doubly special relativity: A kinematics of quantum gravity? // arXiv.org/abs/hep-th/0209264;
Lukierski J., and Nowicki A. Four classes of modified relativistic symmetry transformations // arXiv.org/abs/hep-th/0210111.
 25. *Ahluwalia D. V., Kirchbach M., and Dadhich N.* Operational indistinguishability of doubly special relativities from special relativity // arXiv.org/abs/gr-qc/0212128.

Received 1.13.2003
 Accepted 3.3.2003

ЗАУВАЖЕННЯ ПРО ТРИВІАЛЬНІСТЬ κ -ДЕФОРМАЦІЙ ГРАВІТАЦІЇ

Д. Грумілер, В. Куммер, Д.В. Василевич

Резюме

В рамках $2D$ дилатонної гравітації, сформульованої в термінах пуассонівської сигма-моделі, обговорюються прості спостереження стосовно певних κ -деформацій, увага зосереджується наколо (не-) тривіальності цих деформацій.

ЗАМЕЧАНИЕ О ТРИВИАЛЬНОСТИ κ -ДЕФОРМАЦИЙ ГРАВИТАЦИИ

Д. Грумиллер, В. Куммер, Д.В. Василевич

Резюме

В рамках $2D$ дилатонной гравитации, сформулированной в терминах пуассоновской сигма-модели, обсуждаются простые наблюдения относительно определенных κ -деформаций, вопрос сосредоточен вокруг (не-) тривильности этих деформаций.